

- ment and disability in chronic stroke survivors. *Arch Phys Med Rehabil* 1999;80:1211-8.
19. Green J, Forster A, Bogle S, Young J. Physiotherapy for patients with mobility problems more than 1 year after stroke: a randomized controlled trial. *Lancet* 2002;359:199-203.
 20. Dean CM, Richards CL, Malouin F. Task-related circuit training improves performance of locomotor tasks in chronic stroke: a randomized, controlled pilot trial. *Arch Phys Med Rehabil* 2000;81:409-17.
 21. Hetu FE, Christie CA, Faigenbaum AD. Effects of conditioning on physical fitness and clubhead speed in mature golfers. *Percept Mot Skills* 1998;86:811-5.
 22. Watkins RG, Uppal GS, Perry J, Pink M, Dinsay JM. Dynamic electromyographic analysis of trunk musculature in professional golfers. *Am J Sports Med* 1996;24:535-8.
 23. Pink MM, Jobe FW, Yocum LA, Mottram R. Preventative exercises in golf. *Clin Sports Med* 1996;15:147-61.
 24. Westcott WL, Dolan F, Cavicchi T. Golf and strength training are compatible activities. *Strength Cond* 1996;18:54-6.
 25. Werner R, Kessler S. Effectiveness of an outpatient rehabilitation program for postacute stroke patients. *Am J Phys Med Rehabil* 1996;75:114-20.
 26. Gowland C, Stratford P, Ward M, Moreland J, Torresin W, Van Hullenaar S, et al. Measuring physical impairment and disability with the Chedoke-McMaster Stroke Assessment. *Stroke* 1993;24:58-63.
 27. Schmitz TJ. Sensory assessment. In: O'Sullivan SB, Schmitz TJ, editors. *Physical rehabilitation: assessment and treatment*. 2nd ed. Philadelphia: FA Davis Co; 1988. p. 79-92.
 28. Lamm-Warburg C. Assessment and treatment planning strategies for perceptual deficits. In: O'Sullivan SB, Schmitz TJ, editors. *Physical rehabilitation: assessment and treatment*. 2nd ed. Philadelphia: FA Davis Co; 1988. p. 93-120.
 29. Harburn KL, Hill KM, Kramer JF, Noh S, Vandervoort AA, Matheson JE. Comparison of balance responses to an external perturbation test, with and without an overhead harness safety system. *Gait Posture* 1994;2:27-31.
 30. Cooper MAJ, Mather JSB. Categorisation of golf swings. In: Cochran AJ, Farrally MR, editors. *Science and golf II: Proceedings of the 1994 World Scientific Congress of Golf*. London, UK: E. & F. N. Spon; 1994. p. 65-70.
 31. Robinson RL, Coeur d'Alene A. A study of the correlation between swing characteristics and clubhead velocity. In: Cochran AJ, Farrally MR, editors. *Science and golf II: Proceedings of the 1994 World Scientific Congress of Golf*. London, UK: E. & F. N. Spon; 1994. p. 84-90.
 32. Watanabe K, Kuroki S, Hokari M, Nishizawa S. Golf swing and skill. In: Farrally MR, Cochran AJ, editors. *Science and golf III: Proceedings of the 1998 World Scientific Congress of Golf*. London: E. & F. N. Spon; 1998. p. 29-39.
 33. Geurts ACH, Nienhuis B, Mulder TW. Intrasubject variability of selected force-platform parameters in the quantification of postural control. *Arch Phys Med Rehabil* 1993;74:1144-50.
 34. Goldie PA, Bach TM, Evans OM. Force platform measures for evaluating postural control: reliability and validity. *Arch Phys Med Rehabil* 1989;70:510-7.
 35. Peroto AO. *Anatomical guide for the electromyographer, the limbs and trunk*. Springfield, IL: Charles C. Thomas Publisher; 1994.
 36. Garland SJ, Willems DA, Ivanova TD, Miller KJ. Recovery of standing balance and functional mobility after stroke. *Arch Phys Med Rehabil* 2003;84:1753-9.
 37. Lee WA. Anticipatory control of postural and task muscles during rapid arm flexion. *J Mot Behav* 1980;12:185-96.
 38. Stevenson TJ, Garland SJ. Standing balance during internally produced perturbation in subjects with hemiplegia: validation of the balance scale. *Arch Phys Med Rehabil* 1996;77:656-62.
 39. Berg KO, Wood-Dauphinee SL, Williams JJ, Maki B. Measuring balance in the elderly: validation of an instrument. *Can J Public Health* 1992;83 Suppl 2:S7-11.
 40. Berg K, Wood-Dauphinee S, Williams JJ. The Balance Scale: reliability assessment with elderly residents and patients with an acute stroke. *Scand J Rehabil Med* 1995;27:27-36.
 41. Gowland C, VanHullenaar S, Torresin W, Moreland J, Vanspall B, Barrecca S, et al. *Chedoke-McMaster Stroke Assessment: development, validation and administration manual*. Hamilton, ON: Chedoke-McMaster Hospitals and McMaster University; 1995.
 42. McDowell I, Newell C. *Measuring health: a guide to rating scales and questionnaires*. New York: Oxford University Press; 1987.
 43. de Bruin AF, de Witte LP, Stevens F, Diederiks JP. *Sickness Impact Profile: the state of the art of a generic functional status measure*. *Soc Sci Med* 1992;35:1003-14.
 44. Jones D. The effects of proprioceptive neuromuscular facilitation flexibility training on the clubhead speed of recreational golfers. In: Farrally MR, Cochran AJ, editors. *Science and golf III: Proceedings of the 1998 World Scientific Congress of Golf*. London, UK: E. & F. N. Spon; 1998. p. 46-50.
 45. Grant T, Brouwer B, Culham E, Vandervoort A. Balance retraining following acute stroke: a comparison of two methods. *Can J Occup Ther* 1997;11:69-73.
 46. Petersen H, Magnusson M, Johansson R, Fransson PA. Auditory feedback of perturbed stance in stroke patients. *Scand J Rehabil Med* 1996;28:217-23.
 47. Drummond AER, Walker MF. A randomized controlled trial of leisure rehabilitation after stroke. *Clin Rehabil* 1995;9:283-90.
 48. Kim P, Warren S, Madill H, Hadley M. Quality of life of stroke survivors. *Qual Life Res* 1999;8:293-301.
 49. Lai SM, Studenski S, Duncan PW, Perera S. Persisting consequences of stroke measured by the Stroke Impact Scale. *Stroke* 2002;33:1840-4.
 50. Newall JT, Wood VA, Hower RL, Tinson DJ. Development of a neurological rehabilitation environment: an observational study. *Clin Rehabil* 1997;11:146-55.
 51. van Straten A, de haan RJ, Limburg M, van den Bos GAM. Clinical meaning of the Stroke-Adapted Sickness Impact Profile-30 and the Sickness Impact Profile-136. *Stroke* 2000;31:2610-5.
 52. Morehouse CA. The super senior golfer. In: Cochran AJ, editor. *Science and golf: Proceedings of the First World Scientific Congress of Golf*. London, UK: E. & F.N. Spon; 1990. p. 14-24.
 53. Potempa K, Braun LT, Tinkell T, Popovich J. Benefits of aerobic exercise after stroke. *Sports Med* 1996;21:337-46.
 54. Lindsay D, Strother R, Walker D. *Golf health, science & rehabilitation*. Calgary, AB: FitforeGolf Inc.; 1997.
 55. Chappuis JL, Johnson GD, Murphy K. The "super six"

- strengtheners for golfers. *Physician Sportsmed* 1996;24: 87-8.
56. Pink M, Perry J, Jobe FW. Electromyographic analysis of the trunk in golfers. *Am J Sports Med* 1993;21:385-8.
 57. Kao JT, Pink M, Jobe FW, Perry J. Electromyographic analysis of the scapular muscles during a golf swing. *Am J Sports Med* 1995;23:19-23.
 58. Shatil S, Garland SJ. Strengthening in a therapeutic golf program for individuals following stroke. *Top Geriatr Rehabil* 2000;15:83-94.
 59. Lemak LJ, Fleisig GS, Welch CM, Martin B, Zvijac JE. Usefulness of partial swings in the rehabilitation of golfers. In: Cochran AJ, Farrally MR, editors. *Science and golf II: Proceedings of the 1994 World Scientific Congress of Golf*. London, UK: E. & F. N. Spon; 1994. p. 14-9.
 60. Lindsay D, Strother R, Walker D, Cuskelly G, Schubert G, Paley B. *FitforeGolf physical therapy level II*. Calgary, AB: FitforeGolf Inc.; 1998.
 61. McTeigue M, Lamb SR, Mottram R, Pirozzolo F. Spine and hip motion analysis during the golf swing. In: Cochran AJ, Farrally MR, editors. *Science and golf II: Proceedings of the 1994 World Scientific Congress of Golf*. London, UK: E. & F. N. Spon; 1994. p. 50-8.
 62. Adlington GS. Proper swing technique and biomechanics of golf. *Clin Sports Med* 1996;15:9-26.
 63. Rogers MW, Hedman LD, Pai Y. Kinetic analysis of dynamic transitions in stance support accompanying voluntary leg flexion movements in hemiparetic adults. *Arch Phys Med Rehabil* 1993;74:19-25.
 64. Patten C. Reeducating muscle force control in older persons through strength training. *Top Geriatr Rehabil* 2000;15:47-59.

APPENDIX

Description of Therapeutic Golf Rehabilitation Program

The six components of TGR include general fitness, motor control and muscle strength, flexibility, speed of movement, balance and postural control, and task- (golf) specific training (see Figure 1).

General fitness: Cardiovascular conditioning is recommended as part of golf fitness programs.^{23,52} Following stroke, cardiovascular function and exercise tolerance are often reduced and can result in as much disability as hemiparesis⁵³; hence, individuals poststroke may require cardiovascular training to improve their exercise tolerance. Warming up is also one of the most important factors in enabling peak performance and preventing injuries associated with golf.⁵⁴ TGR includes stretching for muscles of the neck, rotator cuff, shoulders, wrists, fingers, chest, back and trunk, groin, hamstrings, quadriceps, and calves.

Motor control and muscle strength: Golf strengthening recommendations have been proposed for the general public based on the muscular activity required to golf and injury prevention.^{22,55-57} A detailed description of the

motor control and muscle strength component of TGR has been reported elsewhere.³¹

Flexibility: Ranges of motion needed during the golf swing have been reported.^{59,60} If any or all of these movements are limited by soft tissue changes secondary to long-standing hemiparesis, the golf swing will be affected. Subjects were prescribed individual stretching exercises with the goal of increasing their flexibility.

Speed of movement: Not only does the golf swing require coordinated movement of many joint segments through their ranges of motion, but it happens very fast, with a total swing time (defined as time of initial club movement to time that the ball leaves the clubface) of less than 1.3 seconds.^{59,61} Persons with high amounts of muscle spasticity in either of their upper extremities may demonstrate difficulty producing the speed and trajectory needed for successful ball flight. Addressing this problem often requires swing modification, such as using a one-armed swing to prevent frustration and encourage success with ball contact in the golf task.

Balance and postural control: Proper positioning and postural alignment in each of the swing phases are essential to healthy and successful golf.^{54,60,62} Lindsay and colleagues suggested that decreased balance plays a role in the golf weight transfer in that ineffective or incomplete weight transference may decrease ball distance and lead to biomechanical compensations or injuries.⁶⁰ Given that dynamic stability is decreased in persons following stroke,⁶³ one may expect altered weight transfer patterns during the golf swing in this population. All participants in TGR are trained in each of the following balance tasks: assuming a proper address position, weight shifting bilaterally while in a golf stance, adding trunk rotation to the weight shifts, further adding upper extremity perturbations that mimic the back swing and follow-through golf movements, and maintaining the end of the follow-through position. Balance in golf is further challenged by encouraging increased upper extremity movement speeds, performing the task with eyes closed, and adding swing drills that require coordinated physical strategies in a task-specific manner.

Task- (golf) specific training: Practicing functional tasks is essential to motor learning.⁶⁴ Practicing golf is essential to improving one's game.^{58,62} Participants in TGR have six standardized golf lessons, once per week for each of the 6 weeks of the program. For weeks 1 to 3, the lessons focussed on driving, woods, and iron shots; week 4 focussed on irons and chipping; and week 5 focussed on chipping and putting; subjects played nine holes at an accessible golf course in week 6. Grip and swing mechanics are modified according to the participants' stage of motor recovery and golf limitations or problems. Some examples of grip modifications include use of a bilateral baseball grip, a flexible wrist splint, or a modified golf glove that maintains the paretic hand on the club.